arXiv:math/0702601v2 [math.MG] 21 Sep 2007

Rigidity and Volume Preserving Deformation on Degenerate Simplicies

Lizhao Zhang [*]

Abstract

Given a degenerate $(n+1)$-simplex in a Euclidean space R^n and a k with $1 \leq k \leq n$, we separate all its k-faces into 2 groups by following certain rules. The vertices are allowed to have continuous motion in R^n while the volume of the k-faces in the 1st group can not increase (these faces are called "k-cables"), and the volume of the k-faces in the 2nd group can not decrease ("k-struts"). Assuming some smoothness property, we prove that all the volumes of the k-faces will be preserved for any sufficiently small motion. When the vertices are allowed to move in R^{n+1}, we derive a n-degree "characteristic polynomial" for the points configuration, and prove that this property still holds if the $(k-1)$-th coefficient of the polynomial has the desired sign. We also show that all the roots of the polynomial are real.

We generalize the above results to spherical space S^n and hyperbolic space H^n as well.

1 Introduction

Rigidity of geometric structures started to draw research interest from the old times. One of the first substantial mathematical results concerning rigidity is Cauchy's rigidity theorem [2], which says that "Two convex polyhedra comprised of the same number of equal similarly placed faces are superposable or symmetric." When the convexity restriction is removed, a longstanding conjecture (mentioned by Euler) was that there is no flexible polyhedron. Nevertheless a counterexample was found by Connelly [5]. Sabitov [9] later proved that, if a flexible polyhedron is homeomorphic to a sphere, then the volume bounded by the surfaces is constant during the flex. For general polyhedral surface, it was proved by Connelly, Sabitov, and Walz in "The bellows conjecture" [6].

In this paper, our focus is different from the above works in two aspects. First, for a geometric structure, instead of focusing on distance restrictions between some vertices, we analyze the volume restrictions on the k-faces. There are discussions of this type of rigidity in Tay, White and Whiteley([10],[11]). Second, besides Euclidean space, we generalize our results to spherical space and hyperbolic space as well.

---

[*] A main part of the research was conducted at Massachusetts Institute of Technology

In section 2, we study rigidity properties in Euclidean space. Given a degenerate $(n+1)$-simplex in R^n, the vertices are allowed to have continuous motion, either in R^{n+1} or restricted in R^n. For each k with $1 \leq k \leq n$, we separate all its k-faces into 2 groups. During the motion, it is restricted that the volume of the k-faces in the 1st group can not increase, while the volume of the k-faces in the 2nd group can not decrease. We prove that if the continuous motion is restricted in R^n and is C^1, then all the volumes of its k-faces are preserved for any sufficiently small motion, which we call it *k-unyielding* in R^n (Theorem 2.1). When the continuous motion is in R^{n+1} and real analytic, we derive a sequence of constants $c_0, \ldots, c_n$ (Lemma 2.2). We prove that if c_{k-1} has the desired sign, then it is k-unyielding in R^{n+1}, and its vertices will stay in a common n-dim hyperplane (Theorem 2.2). We also prove that $c_1 = 0$ if and only if all the vertices are on a common sphere in R^n (Corollary 2.2). The main results in this section are Theorem 2.1 and Theorem 2.2.

Section 3 is to study the relationship between these k-unyielding properties by analyzing $c_0, \ldots, c_n$ together. We define a *characteristic polynomial* to be

$$f(x) = \sum_{i=0}^{n} (-1)^i c_i x^{n-i}$$

We prove that all roots of $f(x)$ are real numbers, and give a way to count the number of positive roots (Theorem 3.1). We also prove that, the necessary and sufficient condition for $f(x)$ to have n-repeated roots is that any two edges without a common vertex are perpendicular (Theorem 3.2).

In section 4 we generalize these rigidity properties to spherical space S^n and hyperbolic space H^n. For most results in section 2, we find analogues in section 4 as well. In the computation of the volume of the simplices in S^n and H^n, we find an interesting application of the Schläfli differential formula. Some remarks of the history of the Schläfli differential formula can be found in J. Milnor's paper [8]. Like in the Euclidean space, we derive a sequence of constants $c_0, \ldots, c_n$ (Lemma 4.2), and define a characteristic polynomial $f(x)$. We conjecture that $f(x)$ has n real roots in the H^n case (Conjecture 4.1). We prove that if c_{k-1} has the desired sign, then the framework is k-unyielding in S^{n+1} (or H^{n+1}), and the vertices will stay in a common S^n (or H^n) (Theorem 4.4). We also prove that $c_1 = 0$ if and only if all the vertices are in a common n-dim hyperplane (Corollary 4.1).

For the 1-dim hyperbolic space H^1, we derive an inner product space structure (Theorem 4.5), where Cauchy's determinant identity [3] plays a role in proving it. We conjecture that we can derive an inner product space structure for H^n with general n as well (Conjecture 4.2). The main results in this section are Theorem 4.3 and Theorem 4.4.

Since R^n, S^n and H^n are spaces with constant sectional curvature 0, 1 and -1 respectively, we hope this work can help us to find more common rigidity properties among these spaces. This paper is based on and is an extension of the author's PhD thesis [12].

Acknowledgment: The author would like to thank Professor D. Kleitman, R. Stanley, R. Connelly, Wei Luo, and Xun Dong for their many helpful suggestions and discussions on parts of this paper.

2 Rigidity and volume preserving deformation in R^n

2.1 Definition

Given a degenerate $(n+1)$-simplex in Euclidean space, the vertices are allowed to have continuous motion under certain restriction on the volumes of its k-faces.

A k-face is called a *k-cable* if its volume can not increase, a *k-strut* if its volume can not decrease, and a *k-bar* if its volume can not change. We call a framework to be a *k-tensegrity framework* if some of its k-faces are labeled as either k-cables, k-struts, or k-bars. For a framework in R^d, if all the volumes of its k-faces are preserved for any sufficiently small continuous motion under the volume restriction, then we say that it is *k-unyielding* in R^d.

For a framework in R^d, we say that it is *rigid* in R^d, if the distance between each pair of vertices can not be changed for any continuous motion under the volume restriction; and we say that it is *globally rigid* in R^d if the "continuous" restriction is removed.

Notice that, "globally rigid" implies "rigid", which also implies "k-unyielding", while not necessarily the other way around. Some references to the work on 1-tensegrity frameworks can be found in Connelly [4].

2.2 Construction of k-tensegrity frameworks

Given a degenerate $(n+1)$-simplex in R^n with the vertices $A_1, \ldots, A_{n+2}$ in general position (which means that every $n+1$ points are not in a $(n-1)$-dim hyperplane). Easy to see that there is a unique sequence of non-zero coefficients $\alpha_1, \ldots, \alpha_{n+2}$ (up to a non-zero factor), such that $\sum \alpha_i = 0$ and $\sum \alpha_i A_i = 0$. We will use these coefficients throughout this paper.

We choose α_1 to be positive, and separate these $n+2$ points into 2 sets X_1 and X_2 by the following rule. A_i is in X_1 if α_i is positive, and in X_2 if α_i is negative. The separation of these points follows *Radon's theorem*, which says that every $n+2$ points in R^n can be separated into 2 sets such that the convex hulls of the 2 sets have a non-empty intersection. Given k with $1 \leq k \leq n$, we separate all the k-faces into 2 sets $Y_{k,1}$ and $Y_{k,2}$ by the following rule. A k-face is in $Y_{k,1}$ if it has odd number of vertices in X_2, and in $Y_{k,2}$ if it has even number of vertices in X_2. Based on the above separation of k-faces, we construct 2 different k-tensegrity frameworks below.

Framework $G_{n,k}$: let all the k-faces in group $Y_{k,1}$ be k-cables, and all the k-faces in group $Y_{k,2}$ be k-struts.
Framework $F_{n,k}$: let all the k-faces in group $Y_{k,1}$ be k-struts, and all the k-faces in group $Y_{k,2}$ be k-cables.

Notice that $F_{n,k}$ is constructed by switching the role of k-cable and k-strut in $G_{n,k}$. The main purpose of this section is to see whether $G_{n,k}$ and $F_{n,k}$ are k-unyielding in R^n and R^{n+1}.

For $k = 1$, Bezdek and Connelly [1] proved that $G_{n,1}$ is globally rigid in R^{n+1}, and $F_{n,1}$ is rigid in R^n. Besides $G_{n,1}$ and $F_{n,1}$, any other no-bar 1-tensegrity framework constructed on points $A_1, \ldots, A_{n+2}$ is not rigid in R^n; and $G_{n,1}$ is also the only no-bar 1-tensegrity framework to be rigid in R^{n+1}.

The above rigidity properties of $G_{n,1}$ and $F_{n,1}$ are determined solely by how points A_1, …, A_{n+2} are separated into 2 groups. However, for $G_{n,k}$ and $F_{n,k}$ with a general k, we will show that their k-unyielding properties are also determined by other factors.

For $k = n$, easy to see that both $G_{n,n}$ and $F_{n,n}$ are n-unyielding in R^n. The sum of the volume of the n-cables equals the sum of the volume of the n-struts, which is the volume of the convex hull of the $n+2$ points, so non of the volume can change. $G_{n,n}$ and $F_{n,n}$ are not rigid in R^n though, as they can have continuous affine motion while keeping all the volume of their n-faces unchanged.

A natural question to ask is that whether $G_{n,n}$ and $F_{n,n}$ are still n-unyielding in R^{n+1}. We will give the relatively more complicated answer in later sections.

2.3 Exterior algebra

We briefly go over some properties of exterior algebra, which will be used to handle the computation of volumes.

Let $\Lambda(R^n)$ denote the exterior algebra on R^n. If a and b are two elements of $\Lambda(R^n)$, we will denote the exterior product of a and b by $a \wedge b$. A *decomposable k-vector* is an exterior product of k elements of R^n, and a *k-vector* is a linear combination of decomposable k-vectors. Denote $\Lambda^k(R^n)$ the vector space that contains all the k-vectors. We have $\Lambda(R^n) = \bigoplus_{k=0}^{n} \Lambda^k(R^n)$ as a vector space.

If P_1, …, P_{k+1} are $k+1$ points in R^n, we define $\overrightarrow{P_1 \cdots P_{k+1}}$ to be a k-vector as

$$\begin{aligned}\overrightarrow{P_1 \cdots P_{k+1}} &:= \overrightarrow{P_1P_2} \wedge \cdots \wedge \overrightarrow{P_1P_{k+1}} \\ &= \sum_{i=1}^{k+1} (-1)^{i+1} P_1 \wedge \cdots \wedge \hat{P_i} \wedge \cdots \wedge P_{k+1}\end{aligned}$$

Based on the inner product defined on R^n, the inner product on $\Lambda^k(R^n)$ can be well defined as

$$(r_1 \wedge \cdots \wedge r_k) \cdot (s_1 \wedge \cdots \wedge s_k) := \det(r_i \cdot s_j)_{1 \le i,j \le k}$$

where r_1, …, r_k, s_1, …, s_k are $2k$ elements in R^n.

Denote $V_k(P_1, \ldots, P_{k+1})$ the volume of the k-simplex with vertices P_1, …, P_{k+1}. Easy to see that

$$\overrightarrow{P_1 \cdots P_{k+1}}^2 = (k!)^2 V_k^2(P_1, \ldots, P_{k+1})$$

and if we use $l_{i,j}$ to denote $|P_i - P_j|$, we have

$$\begin{aligned}\overrightarrow{P_1 \cdots P_{k+1}}^2 &= (\overrightarrow{P_1P_2} \wedge \cdots \wedge \overrightarrow{P_1P_{k+1}})^2 \\ &= \det(\overrightarrow{P_1P_i} \cdot \overrightarrow{P_1P_j})_{2 \le i,j \le k+1} \\ &= \det((l_{1,i}^2 + l_{1,j}^2 - l_{i,j}^2)/2)_{2 \le i,j \le k+1}\end{aligned} \tag{2.1}$$

We will write the partial derivative of $\overrightarrow{P_1 \cdots P_{k+1}}^2$ over variable $l_{i,j}^2$ as $\partial_{l_{i,j}^2} \overrightarrow{P_1 \cdots P_{k+1}}^2$, and we find that it can be represented as the inner product of two $(k-1)$-vectors, which is more convenient to be used in computation than its polynomial form.

Lemma 2.1

$$\partial_{l^2_{i,j}} \overrightarrow{P_1 \cdots P_{k+1}}^2 = \overrightarrow{P_i P_1 \cdots \hat{P_i} \cdots \hat{P_j} \cdots P_{k+1}} \cdot \overrightarrow{P_j P_1 \cdots \hat{P_i} \cdots \hat{P_j} \cdots P_{k+1}}$$

Proof. By symmetry, we only need to prove the case when $1 < i$ and $1 < j$. Let $A = ((l^2_{1,i} + l^2_{1,j} - l^2_{i,j})/2)_{2 \le i,j \le k+1}$. By formula (2.1), we have $\overrightarrow{P_1 \cdots P_{k+1}}^2 = \det(A)$.

Since $l^2_{i,j}$ only appears inside the $(i-1, j-1)$-th and $(j-1, i-1)$-th term of matrix A, and A is symmetric, so $\partial_{l^2_{i,j}} \det(A)$ is $(-1)^{i+j-1}$ times the determinant of the $(k-1) \times (k-1)$ submatrix of A without the $(i-1)$-th row and $(j-1)$-th column. So

$$\begin{aligned} &\partial_{l^2_{i,j}} \det(A) \\ =\ &(-1)^{i+j-1} (\overrightarrow{P_1P_2} \wedge \cdots \wedge \widehat{\overrightarrow{P_1P_i}} \wedge \cdots \wedge \overrightarrow{P_1P_{k+1}}) \cdot (\overrightarrow{P_1P_2} \wedge \cdots \wedge \widehat{\overrightarrow{P_1P_j}} \wedge \cdots \wedge \overrightarrow{P_1P_{k+1}}) \\ =\ &\overrightarrow{P_i P_1 \cdots \hat{P_i} \cdots \hat{P_j} \cdots P_{k+1}} \cdot \overrightarrow{P_j P_1 \cdots \hat{P_i} \cdots \hat{P_j} \cdots P_{k+1}}. \end{aligned}$$

□

Now we consider the continuous motion $P_i(t)$ with $t \ge 0$ and $P_i(0) = P_i$. To avoid writing a formula too wide in a line, we use p_i to replace $P_i(t)$ to save space.

Corollary 2.1 *If $P_1(t)$, …, $P_{k+1}(t)$ are C^1 and denote $l_{i,j}(t) = |P_i(t) - P_j(t)|$, then*

$$(\overrightarrow{p_1 \cdots p_{k+1}}^2)' = \sum_{i<j} (\overrightarrow{p_i p_1 \cdots \hat{p_i} \cdots \hat{p_j} \cdots p_{k+1}} \cdot \overrightarrow{p_j p_1 \cdots \hat{p_i} \cdots \hat{p_j} \cdots p_{k+1}}) (l^2_{i,j}(t))'$$

Proof. Use chain rule and Lemma 2.1. □

Lemma 2.2 *Embed R^n into a bigger space R^d, and let P and Q are any 2 points. Then*

$$c_k := \sum_{i_1 < \cdots < i_k} \alpha_{i_1} \cdots \alpha_{i_k} \overrightarrow{P A_{i_1} \cdots A_{i_k}} \cdot \overrightarrow{Q A_{i_1} \cdots A_{i_k}}$$

is independent of the choice of points P and Q.

Proof. We prove a stronger result that, for any two l-vectors ω_1 and ω_2 in R^d,

$$\begin{aligned} &c_k(\omega_1, \omega_2; P_1, \ldots, P_k, Q_1, \ldots, Q_k) \\ := &\frac{1}{k!} \sum_{i_1} \cdots \sum_{i_k} \alpha_{i_1} \cdots \alpha_{i_k} (\omega_1 \wedge \overrightarrow{P_1 A_{i_1}} \wedge \cdots \wedge \overrightarrow{P_k A_{i_k}}) \cdot (\omega_2 \wedge \overrightarrow{Q_1 A_{i_1}} \wedge \cdots \wedge \overrightarrow{Q_k A_{i_k}}) \end{aligned}$$

is independent of points P_1,…,P_k,Q_1,…,Q_k.

By symmetry, we only need to prove that it is independent of points P_k and Q_k. Then the problem is reduced to the case of $k = 1$ only.

For $k = 1$, we have

$$\begin{aligned}
&c_1(\omega_1, \omega_2; P, Q)\\
&= \sum_i \alpha_i(\omega_1 \wedge \overrightarrow{PA_i}) \cdot (\omega_2 \wedge \overrightarrow{QA_i})\\
&= \sum_i \alpha_i(\omega_1 \wedge A_i - \omega_1 \wedge P) \cdot (\omega_2 \wedge A_i - \omega_2 \wedge Q)\\
&= \sum_i \alpha_i(\omega_1 \wedge A_i) \cdot (\omega_2 \wedge A_i) + \sum_i \alpha_i(\omega_1 \wedge P) \cdot (\omega_2 \wedge Q)\\
&\quad - (\omega_1 \wedge P) \cdot (\omega_2 \wedge \sum_i \alpha_i A_i) - (\omega_1 \wedge \sum_i \alpha_i A_i) \cdot (\omega_2 \wedge Q)\\
&= \sum_i \alpha_i(\omega_1 \wedge A_i) \cdot (\omega_2 \wedge A_i),
\end{aligned}$$

which is independent of points P and Q. □

2.4 Main theorems

Denote $\mathbf{A}(t) = (A_1(t), \ldots, A_{n+2}(t))$ the continuous motion of points A_1, …, A_{n+2} with $t \geq 0$ and $A_i(0) = A_i$. When $\mathbf{A}(t)$ is restricted in R^n, easy to that there are functions $\alpha_1(t)$, …, $\alpha_{n+2}(t)$ and $h(t)$ that satisfy $\sum \alpha_i(t) = 0$, $\sum \alpha_i(t)A_i(t) = 0$, and

$$|\alpha_i(t)| = h(t) \cdot |\overrightarrow{A_1(t) \cdots \hat{A_i}(t) \cdots A_{n+2}(t)}|,$$

where $h(t)$ is independent of i. When t is small enough, $\alpha_i(t)$ has the same sign as $\alpha_i(0) = \alpha_i$.

Similar to the definition of c_k in Lemma 2.2, we define $c_k(t)$ as well. To avoid writing a formula too long in a line, we use a_i to replace $A_i(t)$ to save space.

Lemma 2.3 *If $\mathbf{A}(t)$ is restricted in R^n and is C^1, then we have*

$$\sum_{i_1 < \cdots < i_{k+1}} \alpha_{i_1}(t) \cdots \alpha_{i_{k+1}}(t)(\overrightarrow{a_{i_1} \cdots a_{i_{k+1}}}^2)' = 0.$$

Proof. For $k = 1$, we have

$$\begin{aligned}
&\sum_{i_1 < i_2} \alpha_{i_1}(t)\alpha_{i_2}(t)(\overrightarrow{a_{i_1}a_{i_2}}^2)'\\
&= -h^2(t) \sum_{i_1 < i_2} (\overrightarrow{a_{i_1}a_1 \cdots \hat{a_{i_1}} \cdots \hat{a_{i_2}} \cdots a_{n+2}} \cdot \overrightarrow{a_{i_2}a_1 \cdots \hat{a_{i_1}} \cdots \hat{a_{i_2}} \cdots a_{n+2}})(\overrightarrow{a_{i_1}a_{i_2}}^2)'
\end{aligned}$$

(by Corollary 2.1)

$$\begin{aligned}
&= -h^2(t)(\overrightarrow{a_1 \cdots a_{n+2}}^2)'\\
&= 0.
\end{aligned}$$

For $k \geq 2$, we have

$$\sum_{i_1 < \cdots < i_{k+1}} \alpha_{i_1}(t) \cdots \alpha_{i_{k+1}}(t) (\overrightarrow{a_{i_1} \cdots a_{i_{k+1}}}^2)'$$

(by Corollary 2.1)

$$\begin{aligned}
&= \sum_{i_1 < \cdots < i_{k+1}} \alpha_{i_1}(t) \cdots \alpha_{i_{k+1}}(t) (\sum_{j<l} (\overrightarrow{a_{i_j} a_{i_1} \cdots \hat{a_{i_j}} \cdots \hat{a_{i_l}} \cdots \ a_{i_{k+1}}} \\
&\quad \cdot \overrightarrow{a_{i_l} a_{i_1} \cdots \hat{a_{i_j}} \cdots \hat{a_{i_l}} \cdots \ a_{i_{k+1}}}) (\overrightarrow{a_{i_j} a_{i_l}}^2)') \\
&= \sum_{j<l} \alpha_j(t) \alpha_l(t) (\overrightarrow{a_j a_l}^2)' \sum_{i_1 < \cdots < i_{k-1}} \alpha_{i_1}(t) \cdots \alpha_{i_{k-1}}(t) \overrightarrow{a_j a_{i_1} \cdots a_{i_{k-1}}} \cdot \overrightarrow{a_l a_{i_1} \cdots a_{i_{k-1}}}
\end{aligned}$$

(by Lemma 2.2)

$$\begin{aligned}
&= \sum_{j<l} \alpha_j(t) \alpha_l(t) (\overrightarrow{a_j a_l}^2)' \cdot c_{k-1}(t) \\
&= 0
\end{aligned}$$

□

Then it leads to the following main theorem.

Theorem 2.1 *If $\mathbf{A}(t)$ is C^1, then $G_{n,k}$ and $F_{n,k}$ are k-unyielding in R^n for each k.*

Proof. For $G_{n,k}$, no matter $A_{i_1} \cdots A_{i_{k+1}}$ is a k-cable or a k-strut, by definition we have

$$\alpha_{i_1}(t) \cdots \alpha_{i_{k+1}}(t) (\overrightarrow{A_{i_1}(t) \cdots A_{i_{k+1}}(t)}^2)' \geq 0.$$

By using Lemma 2.3, we find that

$$(\overrightarrow{A_{i_1}(t) \cdots A_{i_{k+1}}(t)}^2)' = 0$$

must hold for small $t \geq 0$. So $G_{n,k}$ is k-unyielding in R^n. The $F_{n,k}$ case can be proved similarly. □

Remark. For Theorem 2.1, we proved two special cases $k = 1$ and $k = n$ before without using the restriction "$\mathbf{A}(t)$ is C^1". We believe that, $G_{n,k}$ and $F_{n,k}$ can also be proved to be k-unyielding in R^n without using this restriction.

To see that whether $G_{n,k}$ and $F_{n,k}$ are also k-unyielding in R^{n+1}, we will show that the constants $c_0, \ldots, c_n$ we derived in Lemma 2.2 play a main role in it.

Suppose $\mathbf{A}(t)$ moves in R^{n+1}, let $A_0(t)$ be $A_1(t)$'s mirror image over the hyperplane which contains points $A_2(t), \ldots, A_{n+2}(t)$. Easy to see that there are functions $\alpha_1(t), \ldots, \alpha_{n+2}(t)$ that satisfy

$$\frac{1}{2}\alpha_1(t) + \frac{1}{2}\alpha_1(t) + \sum_{i \geq 2} \alpha_i(t) = \sum \alpha_i(t) = 0,$$

and

$$\frac{1}{2}\alpha_1(t)A_0(t) + \frac{1}{2}\alpha_1(t)A_1(t) + \sum_{i\geq 2}\alpha_i(t)A_i(t) = 0.$$

When t is small, $\alpha_i(t)$ has the same sign as $\alpha_i(0) = \alpha_i$

If two functions have the same leading term in the formal power series at $t = 0$, then we will use the symbol "$\approx$".

Lemma 2.4 *Suppose* $\mathbf{A}(t)$ *moves in* R^{n+1} *and is real analytic. If* $c_{k-1} > 0$ *(or* < 0*), then*

$$\sum_{1\leq i_1<\cdots<i_{k+1}} \alpha_{i_1}(t)\cdots\alpha_{i_{k+1}}(t)(\overrightarrow{a_{i_1}\cdots a_{i_{k+1}}}^2)' \leq 0 \quad (or \geq 0)$$

for small $t \geq 0$*, and the equality holds if and only if* $A_1(t)$*, …,* $A_{n+2}(t)$ *stay in a common* n*-dim hyperplane.*

Proof. By symmetry, we only need to prove the case that $c_{k-1} > 0$.

$$\begin{aligned}
&\sum_{1\leq i_1<\cdots<i_{k+1}} \alpha_{i_1}(t)\cdots\alpha_{i_{k+1}}(t)(\overrightarrow{a_{i_1}\cdots a_{i_{k+1}}}^2)' \\
=&\sum_{2\leq i_1<\cdots<i_{k+1}} \alpha_{i_1}(t)\cdots\alpha_{i_{k+1}}(t)(\overrightarrow{a_{i_1}\cdots a_{i_{k+1}}}^2)' \\
&+\frac{1}{2}\alpha_1(t)\sum_{j=1}^{2}\sum_{2\leq i_1<\cdots<i_k} \alpha_{i_1}(t)\cdots\alpha_{i_k}(t)(\overrightarrow{a_j a_{i_1}\cdots a_{i_k}}^2)'
\end{aligned}$$

(by using Lemma 2.3)

$$\begin{aligned}
&= -\frac{1}{2}\alpha_1(t)\cdot\frac{1}{2}\alpha_1(t)\sum_{2\leq i_1<\cdots<i_{k-1}} \alpha_{i_1}(t)\cdots\alpha_{i_{k-1}}(t)(\overrightarrow{a_0 a_1 a_{i_1}\cdots a_{i_{k-1}}}^2)' \\
&\approx -\frac{1}{4}\alpha_1^2(t)\cdot(\overrightarrow{a_0 a_1}^2)'\sum_{2\leq i_1<\cdots<i_{k-1}} \alpha_{i_1}(t)\cdots\alpha_{i_{k-1}}(t)(\overrightarrow{a_0 a_{i_1}\cdots a_{i_{k-1}}}\cdot\overrightarrow{a_1 a_{i_1}\cdots a_{i_{k-1}}}) \\
&\approx -\frac{1}{4}\alpha_1^2(t)\cdot(\overrightarrow{a_0 a_1}^2)'\cdot c_{k-1} \\
&\leq 0
\end{aligned}$$

The equality holds if and only if $A_0(t) = A_1(t)$, namely, $A_1(t)$,…,$A_{n+2}(t)$ stay in a common n-dim hyperplane. □

Then we have the following main theorem.

Theorem 2.2 *Suppose* $\mathbf{A}(t)$ *moves in* R^{n+1} *and is real analytic.*

(1) If $c_{k-1} > 0$*, then* $G_{n,k}$ *is* k*-unyielding in* R^{n+1}*, and points* $A_1(t)$*,…,*$A_{n+2}(t)$ *will stay in a common* n*-dim hyperplane in* R^{n+1} *for small* $t \geq 0$*.*

(2) If $c_{k-1} < 0$*, then* $F_{n,k}$ *is* k*-unyielding in* R^{n+1}*, and points* $A_1(t)$*,…,*$A_{n+2}(t)$ *will stay in a common* n*-dim hyperplane in* R^{n+1} *for small* $t \geq 0$*.*

Proof. We only need to prove (1), as (2) can be proved similarly. Suppose $c_{k-1} > 0$. If points $A_1(t), \ldots, A_{n+2}(t)$ do not stay in a common n-dim hyperplane in R^{n+1} for small $t \geq 0$, then $A_1(t) \neq A_0(t)$. By Lemma 2.4, we have

$$\sum_{1 \leq i_1 < \cdots < i_{k+1}} \alpha_{i_1}(t) \cdots \alpha_{i_{k+1}}(t) (\overrightarrow{a_{i_1} \cdots a_{i_{k+1}}}^2)' < 0.$$

For $G_{n,k}$, however, by definition we have

$$\alpha_{i_1}(t) \cdots \alpha_{i_{k+1}}(t) (\overrightarrow{a_{i_1} \cdots a_{i_{k+1}}}^2)' \geq 0$$

for all the k-faces $A_{i_1} \cdots A_{i_{k+1}}$, which is a contradiction.

As $A_1(t), \ldots, A_{n+2}(t)$ stay in a common n-dim hyperplane, by applying Theorem 2.1, we then have that $G_{n,k}$ is k-unyielding in R^{n+1}. □

Remark. We believe that Theorem 2.2 can be proved without the restriction that "$\mathbf{A}(t)$ is real analytic".

As $c_0 = 1$ is always bigger than 0, Theorem 2.2 gives another explanation that why $G_{n,1}$ is rigid in R^{n+1}. When $k \geq 2$, it is possible that $c_{k-1} = 0$, and we are interested in knowing what the framework looks like. We will give the answer for $k = 2$ next.

2.5 2-tensegrity frameworks $G_{n,2}$ and $F_{n,2}$

Previously we proved that the sign of c_1 determines if $G_{n,2}$ or $F_{n,2}$ is 2-unyielding in R^{n+1}. A natural question to ask is, when does $c_1 = 0$ happen? The answer is amazingly simple, $c_1 = 0$ if and only if points $A_1, \ldots, A_{n+2}$ lie on a common sphere in R^n. Consequently, if we use S_i^{n-1} to denote the sphere in R^n that contains points $A_1, \ldots, \hat{A}_i, \ldots, A_{n+2}$, then $c_1 > 0$ if and only if A_1 is outside the sphere S_1^{n-1}; and $c_1 < 0$ if and only if A_1 is inside the sphere S_1^{n-1}.

Corollary 2.2 *$c_1 = 0$ if and only if A_1 lies on S_1^{n-1}, namely, points $A_1, \ldots, A_{n+2}$ lie on a common sphere in R^n.*

Proof. We define a transformation f in R^n, such that for any point $P \neq A_1$, $f(P)$ is a point that satisfies

$$\overrightarrow{A_1 f(P)} = \overrightarrow{A_1 P} / \overrightarrow{A_1 P}^2.$$

Since $\sum_{i=2}^{n+2} \alpha_i \overrightarrow{A_1 A_i} = 0$, so we have

$$\sum_{i=2}^{n+2} (\alpha_i \overrightarrow{A_1 A_i}^2) \overrightarrow{A_1 f(A_i)} = 0.$$

A basic property in *inversion geometry* is that, A_1 lies on S_1^{n-1} if and only if $f(S_1^{n-1})$ is a $(n-1)$-dim hyperplane. Since $f(S_1^{n-1})$ is a $(n-1)$-dim hyperplane if and only if $\sum_{i=2}^{n+2} \alpha_i \overrightarrow{A_1 A_i}^2 = 0$, which is also equivalent to $c_1 = 0$, so A_1 lies on S_1^{n-1} if and only if $c_1 = 0$. □

To show some geometric features of $G_{n,2}$ and $F_{n,2}$, we use $n = 2$ as example, and *assume* for a moment that Theorem 2.2 is also true without the smoothness requirement. Suppose A_1, A_2, A_3 and A_4 are vertices of a convex quadrilateral in a plane. Topologically, it is not so obvious for us to tell the difference between $G_{2,2}$ and $F_{2,2}$, so how can we determine that which one of them is 2-unyielding in R^3? In Figure 1 (a), A_1 is inside the dotted circle, then Theorem 2.2 tells us that $c_1 < 0$ and $F_{2,2}$ is 2-unyielding in R^3; and $A_1(t)$, …, $A_4(t)$ will stay in a common 2-dim plane in R^3 for small t. In Figure 1 (b), A_1 is outside the dotted circle, then $c_1 > 0$ and $G_{2,2}$ is 2-unyielding in R^3.



Figure 1:

An even more interesting explanation comes with the framework whose four 2-faces are all 2-bars (whose area are fixed in the continuous motion). Theorem 2.2 says that, in order for $A_1(t) \cdots A_4(t)$ to become a non-degenerate 3-simplex, the vertices will have to first stay in a common 2-dim plane until they move on to a common circle, then it can be "freed" to R^3.

It will be interesting to know the geometric meaning of $c_k = 0$ when $k \geq 2$.

3 Characteristic polynomial of points in R^n

3.1 Definition of characteristic polynomial

In the previous section, we studied the k-unyielding properties of $G_{n,k}$ and $F_{n,k}$ for $1 \leq k \leq n$. In this section, we will show some relationship between these k-unyielding properties with different k's. Remember that we derived a sequence of constants c_0, …, c_n in Lemma 2.2, and showed in Theorem 2.2 that the sign of c_{k-1} plays a main role in deciding whether $G_{n,k}$ and $F_{n,k}$ are k-unyielding in R^{n+1}.

We define a *characteristic polynomial* $f(x)$ to be

$$f(x) = c_0 x^n - c_1 x^{n-1} + \cdots + (-1)^i c_i x^{n-i} + \cdots + (-1)^n c_n \tag{3.1}$$

3.2 Properties of characteristic polynomial

We will show that the roots $\{\lambda_1, \ldots, \lambda_n\}$ of $f(x)$ are all real numbers, and give a way to count the number of positive roots.

Let B be a $(n+1)\times n$ matrix whose i-th row is vector $\overrightarrow{A_{n+2}A_i}$, and D be a $(n+1)\times(n+1)$ diagonal matrix whose i-th diagonal term is α_i.

Proposition 3.1 *The characteristic polynomials of matrix BB^TD and B^TDB are $xf(x)$ and $f(x)$ respectively.*

Proof. The coefficient of x^{n+1-k} in the characteristic polynomial of BB^TD is $(-1)^k$ times the sum of the determinants of all $k \times k$ main diagonal submatrices of BB^TD, which is c_k by Lemma 2.2. Therefore $xf(x)$ is the characteristic polynomial of BB^TD, and then $f(x)$ is the characteristic polynomial of B^TDB. $\square$

Theorem 3.1 *All λ_i are non-zero real numbers. If $\{\alpha_1,\ldots,\alpha_{n+2}\}$ have s positive numbers, then $\{\lambda_1,\ldots, \lambda_n\}$ have $s-1$ positive numbers and $n+1-s$ negative numbers.*

Proof. By Proposition 3.1, $f(x)$ is the characteristic polynomial of symmetric matrix B^TDB, so all λ_i are real. Easy to see that B^TDB has full rank, so all λ_i are also non-zero. Therefore $\{\lambda_1,\ldots, \lambda_n\}$ have the same signs as a n-subset of the diagonal terms of D. By symmetry, for any j with $1 \le j \le n+2$, $\{\lambda_1,\ldots, \lambda_n\}$ should have the same signs as a n-subset of $\{\alpha_1, \ldots, \hat{\alpha_j}, \ldots, \alpha_{n+2}\}$. So $\{\lambda_1,\ldots, \lambda_n\}$ must have $s-1$ positive numbers and $n+1-s$ negative numbers. $\square$

Theorem 3.2 *The necessary and sufficient condition for the characteristic polynomial $f(x)$ to have n-repeated roots is that $\overrightarrow{A_iA_j} \cdot \overrightarrow{A_kA_l} = 0$ for all distinct numbers i, j, k and l.*

Proof.

(1) If $\lambda_1 = \cdots = \lambda_n = \lambda$, then by Proposition 3.1, $B^TDB = \lambda I_n$ where I_n is the $n \times n$ identity matrix. Let B_1 be a $(n+1) \times (n+1)$ matrix, whose first n columns are B, and every term on the last column is $\sqrt{-\lambda/\alpha_{n+2}}$. Easy to see that $B_1^TDB_1 = \lambda I_{n+1}$, so $B_1B_1^TD = \lambda I_{n+1}$ as well. Therefore the (i,j)-th term of $B_1B_1^TD$ is 0 if $i \neq j$, which means that
$$\overrightarrow{A_{n+2}A_i} \cdot \overrightarrow{A_{n+2}A_j} = \frac{\lambda}{\alpha_{n+2}}.$$
So if i, j, k and $n+2$ are distinct, then $\overrightarrow{A_{n+2}A_i} \cdot \overrightarrow{A_jA_k} = 0$. By symmetry, we have $\overrightarrow{A_iA_j} \cdot \overrightarrow{A_kA_l} = 0$ for all distinct numbers i, j, k and l.

(2) Suppose $\overrightarrow{A_iA_j} \cdot \overrightarrow{A_kA_l} = 0$ for all distinct numbers i, j, k and l. So for each i with $1 \le i \le n+2$, $\overrightarrow{A_iA_j} \cdot \overrightarrow{A_iA_k}$ is independent of j and k, as long as i, j and k are distinct. We denote it b_i.

Let B_2 be a $n\times n$ matrix whose i-th row is vector $\overrightarrow{A_{n+2}A_i}$, B_3 be a $n\times n$ matrix whose i-th row is vector $\overrightarrow{A_{n+1}A_i}$, and D_1 be a $n \times n$ diagonal matrix whose i-th diagonal term is α_i. By Lemma 2.2, $f(x)$ is the characteristic polynomial of $B_2B_3^TD_1$. Since

$\overrightarrow{A_{n+2}A_i}\cdot\overrightarrow{A_{n+1}A_j} = 0$ for $i \neq j$ with $1 \leq i, j \leq n$, so the non-diagonal terms of $B_2B_3^TD_1$ are all 0. The i-th diagonal term is

$$\alpha_i\overrightarrow{A_{n+2}A_i}\cdot\overrightarrow{A_{n+1}A_i} = \alpha_i b_i.$$

So $\{\lambda_1, \dots, \lambda_n\}$ are $\{\alpha_1 b_1, \dots, \alpha_n b_n\}$. By symmetry, for any i and j with $1 \leq i < j \leq n+2$, $\{\lambda_1, \dots, \lambda_n\}$ are the same as $\{\alpha_1 b_1, \dots, \widehat{\alpha_i b_i}, \dots, \widehat{\alpha_j b_j}, \dots, \alpha_{n+2} b_{n+2}\}$. So all $\alpha_i b_i$ are equal, and therefore $\lambda_1 = \cdots = \lambda_n$.

□

3.3 Generalization of characteristic polynomial

The concept of characteristic polynomial can be defined in a more general sense when the number of points in R^n is more than $n+2$. Let A_1, …, A_m be m ($m \geq n+2$) points in R^n, and α_1,…,α_m be a sequence of coefficients that satisfy $\sum_{i=1}^m \alpha_i = 0$ and $\sum_{i=1}^m \alpha_i A_i = 0$.

A sequence of c_i ($0 \leq i \leq n$) can still be defined as in Lemma 2.2, and we then define the characteristic polynomial $f(x)$ the same way in (3.1).

Theorem 3.3 *All roots of $f(x)$ are real numbers.*

Proof. Let B be a $m \times n$ matrix whose i-th row is the coordinate of point A_i; D be a $m \times m$ diagonal matrix whose i-th term on diagonal is α_i. By slightly modifying Proposition 3.1, we can show that $f(x)$ is the characteristic polynomial of symmetric matrix B^TDB. So all roots of $f(x)$ are real numbers. □

4 Rigidity and volume preserving deformation in S^n and H^n

4.1 Elementary geometry in S^n and H^n

In this section, we study the k-tensegrity frameworks in spherical space S^n and hyperbolic space H^n. Some basic properties of S^n and H^n can be found in Fenchel's book [7].

For hyperbolic space H^n, we define $R^{n,1}$ (not R^{n+1}) as a $(n+1)$-dim linear space. If we use $(x_1, \dots, x_{n+1})$ as the coordinates of $R^{n,1}$, then the equation that H^n satisfies is

$$-x_1^2 + x_2^2 + \cdots + x_{n+1}^2 = -1, \quad \text{with} \quad x_1 > 0.$$

A bilinear product "·" in $R^{n,1}$ is defined as

$$\overrightarrow{OA}\cdot\overrightarrow{OB} = -x_1y_1 + x_2y_2 + \cdots + x_{n+1}y_{n+1},$$

where $(x_1, \dots, x_{n+1})$ and $(y_1, \dots, y_{n+1})$ are the coordinates of A and B. If A is a point in H^n, then $\overrightarrow{OA}^2 = -1$. This operation "·" gives a quadratic form in $R^{n,1}$, whose restriction on the tangent space at any point in H^n is positive definite, and induces a *Riemannian metric* on H^n. S^n is defined as a sphere with radius 1 in an Euclidean space R^{n+1}, and we use S_+^n to denote the open hemisphere. Both S_+^n and H^n are *geodesic convex*, and they have *constant sectional curvature* 1 and -1 respectively. We use $\widetilde{PQ}$ to denote the *geodesic distance* between 2 points P and Q.

The notion "simplex" is defined in H^n as well, and it follows similarly in S^n_+. Suppose P_1, …, P_{k+1} are $k+1$ points in H^n, then all the linear combination $\sum_{i=1}^{k+1}\gamma_i P_i$ with $\gamma_i \geq 0$ span a *cone* in $R^{n,1}$. The intersection of this cone with H^n is called a *hyperbolic k-simplex*, or *H^k-simplex* for convenience. The hyperbolic k-simplex stays in the k-dim *totally geodesic submanifold* in H^n that contains points P_1, …, P_{k+1}. In the S^n_+ case, it is called *spherical k-simplex*, or *S^k_+-simplex* for convenience. Denote $V_k(P_1,\dots,P_{k+1})$ the k-dim volume of the hyperbolic (or spherical) k-simplex under the Riemannian metric. Notice that $V_1(P_1,P_2)=\widetilde{P_1P_2}$.

Similar to the exterior algebra in Euclidean space, a bilinear product "·" in $\Lambda^k(R^{n,1})$ is well defined as

$$(r_1\wedge\cdots\wedge r_k)\cdot(s_1\wedge\cdots\wedge s_k)=\det(r_i\cdot s_j)_{1\leq i,j\leq k},$$

where r_1, …, r_k, s_1, …, s_k are $2k$ elements in $R^{n,1}$.

If P_1, …, P_{k+1} are $k+1$ points in $R^{n,1}$, we define $\overrightarrow{P_1\cdots P_{k+1}}$ in $\Lambda^k(R^{n,1})$ to be

$$\overrightarrow{P_1\cdots P_{k+1}}=\overrightarrow{P_1P_2}\wedge\overrightarrow{P_1P_3}\wedge\cdots\wedge\overrightarrow{P_1P_{k+1}}.$$

We also define

$$|\overrightarrow{OP_1\cdots P_{k+1}}|=|\overrightarrow{OP_1\cdots P_{k+1}}^2|^{1/2}.$$

If A and B are 2 different points in H^n, then $\overrightarrow{AB}^2>0$, and

$$\overrightarrow{AB}^2=2\cosh\widetilde{AB}-2,\quad A\cdot B=-\cosh\widetilde{AB},\quad \overrightarrow{OAB}^2=-\sinh^2\widetilde{AB};$$

if A and B are in S^n_+, then

$$\overrightarrow{AB}^2=2-2\cos\widetilde{AB},\quad A\cdot B=\cos\widetilde{AB},\quad \overrightarrow{OAB}^2=\sin^2\widetilde{AB}.$$

In both cases we have

$$\frac{d(\overrightarrow{AB}^2)}{d\widetilde{AB}}=2|\overrightarrow{OAB}| \tag{4.1}$$

Formula (4.1) means that, when $\widetilde{AB}$ increases (or decreases), $\overrightarrow{AB}^2$ increases (or decreases) as well. If P_1, …, P_{k+1} are $k+1$ points in S^n_+ (or H^n), then $V_k(P_1,\dots,P_{k+1})$ is a function with all $\overrightarrow{P_iP_j}^2$ as the variables. We will write the partial derivative of $V_k(P_1,\dots,P_{k+1})$ over $\overrightarrow{P_iP_j}^2$ as $\partial_{\overrightarrow{P_iP_j}^2}V_k(P_1,\dots,P_{k+1})$.

4.2 Construction of k-tensegrity frameworks in S^n_+ and H^n

Like in the Euclidean space, we can define k-cable, k-strut, k-bar, and k-tensegrity framework in S^n_+ and H^n similarly.

Given a degenerate $(n+1)$-simplex in S^n_+ (or H^n) with the vertices A_1, …, A_{n+2} in general position (which means that every $n+1$ points are not in a lower dimensional space S^{n-1}_+ (or H^{n-1})). Easy to see that there is a unique sequence of non-zero coefficients α_1, …, α_{n+2} (up to a non-zero factor), such that $\sum\alpha_iA_i=0$.

We choose α_1 to be positive, and construct k-tensegrity framework $G_{n,k}$ and $F_{n,k}$ in the same way we did in section 2.2 in Euclidean space. We will show whether $G_{n,k}$ and $F_{n,k}$ are k-unyielding in S^n_+ (or H^n) and S^{n+1}_+ (or H^{n+1}).

For $k = n$, it is easy to see that $G_{n,n}$ and $F_{n,n}$ are n-unyielding in R^n. The sum of the volume of the n-cables equals the sum of the volume of the n-struts, which is the volume of the convex hull of the $n+2$ points, so non of the volume can change.

4.3 Main theorems

Theorem 4.1 *(1) $G_{n,1}$ is globally rigid in S^{n+1}.*

(2) $F_{n,1}$ is rigid in S^n_+.

Proof.

(1) Suppose $B_1, \ldots, B_{n+2}$ are $n+2$ points in S^{n+1} that satisfy the cable-strut restriction set by $G_{n,1}$. Then we have

$$\begin{aligned}
0 &\le (\alpha_1 B_1 + \cdots + \alpha_{n+2} B_{n+2})^2 \\
&= \sum_j \alpha_j \sum_i \alpha_i B_i^2 - \sum_{i<j} \alpha_i \alpha_j (B_i - B_j)^2 \\
&\le \sum_j \alpha_j \sum_i \alpha_i A_i^2 - \sum_{i<j} \alpha_i \alpha_j (A_i - A_j)^2 \\
&= (\alpha_1 A_1 + \cdots + \alpha_{n+2} A_{n+2})^2 \\
&= 0.
\end{aligned}$$

Then $(B_i - B_j)^2 = (A_i - A_j)^2$ must hold, so $G_{n,1}$ is globally rigid in S^{n+1}.

(2) $F_{n,1}$ is constructed by switching the role of cable and strut in $G_{n,1}$, so "$F_{n,1}$ is rigid in S^n_+" is equivalent to "$G_{n,1}$ is rigid in S^n_+", which we just proved in (1).

□



Figure 2:

An example of $G_{2,1}$ in S^2_+ is given in Figure 2. Surprisingly, the proof of Theorem 4.1 can not be directly applied to prove the H^n case. The main difference is that, in $R^{n,1}$, it is possible for $\overrightarrow{OB}^2$ to be negative. We will use a different method to prove the H^n case.

Theorem 4.2 *(1) $G_{n,1}$ is globally rigid in H^{n+1}.*

(2) $F_{n,1}$ is rigid in H^n.

Proof.

(1) Suppose α_1 ,..., $\alpha_m > 0$ and α_{m+1}, ..., $\alpha_{n+2} < 0$, and suppose that B_1, ..., B_{n+2} are $n+2$ points in H^{n+1} that satisfy the cable-strut restrictions set by $G_{n,1}$. Since $\alpha_1, \ldots, \alpha_m > 0$, so there is a $f_1 > 0$ such that $f_1 \sum_{i=1}^{m} \alpha_i B_i$ is a point on H^{n+1}; similarly, there is a $f_2 > 0$ such that $-f_2 \sum_{i=m+1}^{n+2} \alpha_i B_i$ is a point on H^{n+1}. We denote these 2 points D_1 and D_2, and denote $\beta_i = f_1\alpha_i$ if $1 \le i \le m$ and $\beta_i = f_2\alpha_i$ if $m+1 \le i \le n+2$. Then we have

$$\begin{aligned} 0 &\le (D_1 - D_2)^2 \\ &= (\sum \beta_i B_i)^2 \\ &= \sum_j \beta_j \sum_i \beta_i B_i^2 - \sum_{i<j} \beta_i\beta_j (B_i - B_j)^2 \\ &\le \sum_j \beta_j \sum_i \beta_i A_i^2 - \sum_{i<j} \beta_i\beta_j (A_i - A_j)^2 \\ &= (\sum \beta_i A_i)^2 \\ &= (f_1 \sum_{i=1}^{m} \alpha_i A_i + f_2 \sum_{i=m+1}^{n+2} \alpha_i A_i)^2 \end{aligned}$$

(since $\sum \alpha_i A_i = 0$)

$$= ((f_1 - f_2) \sum_{i=1}^{m} \alpha_i A_i)^2$$

(since $\sum_{i=1}^{m} \alpha_i A_i$ is a multiple of a point on H^{n+1}, so $(\sum_{i=1}^{m} \alpha_i A_i)^2 < 0$)

$$\le 0.$$

So $(B_i - B_j)^2 = (A_i - A_j)^2$ must hold, and $G_{n,1}$ is therefore globally rigid.

(2) $F_{n,1}$ is constructed by switching the cable and strut in $G_{n,1}$, so "$F_{n,1}$ is rigid in H^n" is equivalent to "$G_{n,1}$ is rigid in H^n", which we just proved in (1).

□

In the study of the k-unyielding properties of $G_{n,k}$ and $F_{n,k}$ for general k, Schläfli's differential formula plays a central role in the computation of the volume of the S_+^k(or H^k)-simplices. Some remarks on the history of the Schläfli differential formula can be found in J. Milnor's paper [8].

Given a family of k-dim simplices Δ which vary in a differentiable manner in a space with constant sectional curvature $\kappa = 1$ or $\kappa = -1$, Schläfli differential formula states that

$$\kappa \cdot dV_k(\Delta) = \frac{1}{k-1} \sum_F V_{k-2}(F)\, d\theta_F \tag{4.2}$$

where the sum is taken over all $(k-2)$-faces F of Δ, $V_{k-2}(F)$ denotes the volume of face F and θ_F denotes the dihedral angle at face F.

If Δ has vertices P_1, …, P_{k+1}, then we will use $|\overrightarrow{O\Delta}|$ to denote

$$|\overrightarrow{OP_1 \dots P_{k+1}}|.$$

For $(k-2)$-face F, $|\overrightarrow{OF}|$ is defined similarly.

We also define a notation $\langle \cdot | \cdot ; \dots \rangle$ as

$$\langle Q|P_i; P_1, \dots, P_{k+1}\rangle := (-1)^{i-1} \frac{\overrightarrow{OQP_1 \cdots \hat{P}_i \cdots P_{k+1}} \cdot \overrightarrow{OP_1 \cdots P_{k+1}}}{\overrightarrow{OP_1 \cdots P_{k+1}}^2}$$

Lemma 4.1 *Let P, Q, B_1, …, B_k be the $k+2$ vertices of a S_+^{k+1}(or H^{k+1})-simplex Δ, and $\kappa = 1$ (or $\kappa = -1$) be the constant sectional curvature. Define*

$$d(P, Q; B_1, \dots, B_k) := 2 \cdot (k+1)! \, |\overrightarrow{O\Delta}| \, \partial_{\overrightarrow{PQ}^2} V_{k+1}(\Delta),$$

then $d(P, Q) = 1$ and

$$d(P, Q; B_1, \dots, B_k) = \kappa^k (k+1)(k-1)! \sum_F g_F |\overrightarrow{OF}| V_{k-1}(F),$$

where g_F is defined in the following:

(1) If the vertices of F are B_1, …, B_k, then $g_F = 1$.

(2) If the vertices of F are P, B_1, …, $\hat{B}_i$, …, B_k, then

$$g_F = -\langle B_i | P; P, Q, B_1, \dots, \hat{B}_i, \dots, B_k\rangle.$$

Similarly if the vertices of F are Q, B_1, …, $\hat{B}_i$, …, B_k.

(3) If the vertices of F are P, Q, B_1, …, $\hat{B}_i$, …, $\hat{B}_j$, …, B_k, then

$$\begin{aligned} g_F = {} & \langle B_i | P; P, Q, B_1, \dots, \hat{B}_j, \dots, B_k\rangle \cdot \langle B_j | Q; P, Q, B_1, \dots, \hat{B}_i, \dots, \hat{B}_j, \dots, B_k\rangle \\ & + \langle B_j | P; P, Q, B_1, \dots, \hat{B}_i, \dots, B_k\rangle \cdot \langle B_i | Q; P, Q, B_1, \dots, \hat{B}_i, \dots, \hat{B}_j, \dots, B_k\rangle. \end{aligned}$$

Proof. It can be proved by applying Schläfli differential formula (4.2). The proof only involves straight computation of $\partial_{\overrightarrow{PQ}^2} \theta_F$ for all the $(k-1)$-faces F, but is rather lengthy. As it is not a focal point of this paper, we just skip it. □

In order to see that whether $G_{n,k}$ and $F_{n,k}$ are k-unyielding in S_+^n (or H^n) or S_+^{n+1} (or H^{n+1}), we give the following important result, which is the analogue of Lemma 2.2.

Lemma 4.2 *Embed S_+^n(or H^n) into a bigger space S_+^d (or H^d), and let P and Q be any 2 points in S_+^d (or H^d). Then*

$$c_k := \sum_{1 \le i_1 < \cdots < i_k} \alpha_{i_1} \cdots \alpha_{i_k} d(P, Q; A_{i_1}, \dots, A_{i_k})$$

is independent of the choice of points P and Q, and

$$c_k = \kappa^k (k+1)(k-1)! \sum_{1 \le i_1 < \cdots < i_k} \alpha_{i_1} \cdots \alpha_{i_k} |\overrightarrow{OA_{i_1} \cdots A_{i_k}}|\, V_{k-1}(A_{i_1}, \ldots, A_{i_k}) \tag{4.3}$$

Proof. Apply Lemma 4.1. $\square$

Denote $\mathbf{A}(t) = (A_1(t), \ldots, A_{n+2}(t))$ the continuous motion with $t \ge 0$ and $A_i(0) = A_i$. When $\mathbf{A}(t)$ is restricted in S^n_+ (or H^n), easy to prove that there are functions $\alpha_1(t), \ldots, \alpha_{n+2}(t)$ and $h(t)$ that satisfy $\sum \alpha_i(t) A_i(t) = 0$ and

$$|\alpha_i(t)| = h(t) \cdot |\overrightarrow{OA_1(t) \cdots \hat{A_i}(t) \cdots A_{n+2}(t)}|,$$

where $h(t)$ is independent of i. When t is small enough, $\alpha_i(t)$ has the same sign as $\alpha_i(0) = \alpha_i$. Similar to the definition of c_k in Lemma 4.2, we can define $c_k(t)$ as well.

To avoid writing a formula too wide in a line, we use a_i to replace $A_i(t)$ to save space.

Lemma 4.3 *If $\mathbf{A}(t)$ is restricted in S^n_+ (or H^n) and is C^1, then we have*

$$\sum_{1 \le i_1 < \cdots < i_{k+1}} \alpha_{i_1}(t) \cdots \alpha_{i_{k+1}}(t)\, |\overrightarrow{Oa_{i_1} \cdots a_{i_{k+1}}}|\, (V_k(a_{i_1}, \ldots, a_{i_{k+1}}))' = 0$$

Proof. For $k = 1$, we have

$$2 \sum_{i_1 < i_2} \alpha_{i_1}(t) \alpha_{i_2}(t)\, |\overrightarrow{Oa_{i_1} a_{i_2}}|\, (V_1(a_{i_1}, a_{i_2}))'$$

(by formula (4.1))

$$= \sum_{i_1 < i_2} \alpha_{i_1}(t) \alpha_{i_2}(t)\, (\overrightarrow{a_{i_1} a_{i_2}}^2)'$$

("$-$" is for S^n_+ case and "$+$" is for H^n case)

$$= \mp h^2(t) \cdot \sum_{i_1 < i_2} (\overrightarrow{Oa_{i_1} a_1 \cdots \hat{a_{i_1}} \cdots \hat{a_{i_2}} \cdots a_{n+2}} \cdot \overrightarrow{Oa_{i_2} a_1 \cdots \hat{a_{i_1}} \cdots \hat{a_{i_2}} \cdots a_{n+2}}) (\overrightarrow{a_{i_1} a_{i_2}}^2)'$$

(by Corollary 2.1)

$$= \mp h^2(t) \cdot (\overrightarrow{Oa_1 \cdots a_{n+2}}^2)'$$
$$= 0.$$

For $k \ge 2$, we have

$$2 \cdot k! \sum_{1 \le i_1 < \cdots < i_{k+1}} \alpha_{i_1}(t) \cdots \alpha_{i_{k+1}}(t)\, |\overrightarrow{Oa_{i_1} \cdots a_{i_{k+1}}}|\, (V_k(a_{i_1}, \ldots, a_{i_{k+1}}))'$$

$$= 2 \cdot k! \sum_{1 \le i_1 < \cdots < i_{k+1}} \alpha_{i_1}(t) \cdots \alpha_{i_{k+1}}(t)\, |\overrightarrow{Oa_{i_1} \cdots a_{i_{k+1}}}| \sum_{j<l} \partial_{\overrightarrow{a_{i_j} a_{i_l}}^2} V_k(a_{i_1}, \ldots, a_{i_{k+1}}) (\overrightarrow{a_{i_j} a_{i_l}}^2)'$$

$$= 2 \cdot k! \sum_{j<l} \alpha_j(t)\alpha_l(t)(\overrightarrow{a_j a_l}^2)' \, (\sum_{1 \le i_1 < \cdots < i_{k-1}} \alpha_{i_1}(t) \cdots \alpha_{i_{k-1}}(t) \, |\overrightarrow{O a_j a_l a_{i_1} \cdots a_{i_{k-1}}}|$$

$$\cdot \, \partial_{\overrightarrow{a_j a_l}^2} \, V_k(a_j, a_l, a_{i_1}, \ldots, a_{i_{k-1}}))$$

(by Lemma 4.2)

$$= c_{k-1}(t) \sum_{j<l} \alpha_j(t)\alpha_l(t)(\overrightarrow{a_j a_l}^2)'$$

(we just proved $\sum_{j<l} \alpha_j(t)\alpha_l(t)(\overrightarrow{a_j a_l}^2)' = 0$ above)

$$= 0.$$

□

Then it leads to the following main theorem.

Theorem 4.3 *If $\mathbf{A}(t)$ is C^1, then $G_{n,k}$ and $F_{n,k}$ are k-unyielding in S^n_+ (or H^n) for each k.*

Proof. For $G_{n,k}$, for both k-cable and k-strut, by definition we have

$$\alpha_{i_1}(t) \cdots \alpha_{i_{k+1}}(t)(V_k(A_{i_1}(t), \ldots, A_{i_{k+1}}(t))' \geq 0.$$

By using Lemma 4.3, we find that

$$(V_k(A_{i_1}(t), \ldots, A_{i_{k+1}}(t))' = 0$$

must hold for small t. So $G_{n,k}$ is k-unyielding in S^n_+(or H^n). The same proof works for $F_{n,k}$. □

Remark. We believe that the above theorem can be proved without the restriction "$\mathbf{A}(t)$ is C^1".

We now start to discuss that whether $G_{n,k}$ and $F_{n,k}$ are also k-unyielding in S^{n+1} (or H^{n+1}). Suppose $\mathbf{A}(t)$ moves in S^{n+1} (or H^{n+1}), then there is a unique S^n (or H^n) that contains points $A_2(t)$, …, $A_{n+2}(t)$. Let $A_0(t)$ on S^{n+1} (or H^{n+1}) be the mirror image of $A_1(t)$ over hyperplane $OA_2(t) \cdots A_{n+2}(t)$. Easy to see that there are functions $\alpha_1(t)$, …, $\alpha_{n+2}(t)$ that satisfy

$$\frac{1}{2}\alpha_1(t)A_0(t) + \frac{1}{2}\alpha_1(t)A_1(t) + \sum_{i=2}^{n+2} \alpha_i(t)A_i(t) = 0$$

and when t is small, $\alpha_i(t)$ has the same sign as $\alpha_i(0) = \alpha_i$.

Lemma 4.4 *Suppose $\mathbf{A}(t)$ moves in S^{n+1} or H^{n+1} and is real analytic. If $c_{k-1} > 0$ (or < 0), then*

$$\sum_{1\le i_1<\cdots<i_{k+1}} \alpha_{i_1}(t)\cdots\alpha_{i_{k+1}}(t)\,|\overrightarrow{Oa_{i_1}\cdots a_{i_{k+1}}}|\,(V_k(a_{i_1},\ldots,a_{i_{k+1}}))' \le 0 \quad (\mathit{or} \ge 0)$$

for small $t \ge 0$, and the equality holds if and only if $A_1(t)$, …, $A_{n+2}(t)$ stay in a common S^n (or H^n).

Proof. By symmetry, we only need to prove the case when $c_{k-1} > 0$.

$$\begin{aligned}
&\sum_{1\le i_1<\cdots<i_{k+1}} \alpha_{i_1}(t)\cdots\alpha_{i_{k+1}}(t)\,|\overrightarrow{Oa_{i_1}\cdots a_{i_{k+1}}}|\,(V_k(a_{i_1},\ldots,a_{i_{k+1}}))'\\
=&\sum_{2\le i_1<\cdots<i_{k+1}} \alpha_{i_1}(t)\cdots\alpha_{i_{k+1}}(t)\,|\overrightarrow{Oa_{i_1}\cdots a_{i_{k+1}}}|\,(V_k(a_{i_1},\ldots,a_{i_{k+1}}))'\\
&+\frac{1}{2}\alpha_1(t)\sum_{j=1}^{2}\sum_{2\le i_1<\cdots<i_k} \alpha_{i_1}(t)\cdots\alpha_{i_k}(t)\,|\overrightarrow{Oa_ja_{i_1}\cdots a_{i_k}}|\,(V_k(a_j,a_{i_1},\ldots,a_{i_k}))'
\end{aligned}$$

(by using Lemma 4.3)

$$= -\frac{1}{4}\alpha_1^2(t)\sum_{2\le i_1<\cdots<i_{k-1}} \alpha_{i_1}(t)\cdots\alpha_{i_{k-1}}(t)\,|\overrightarrow{Oa_0a_1a_{i_1}\cdots a_{i_{k-1}}}|\,(V_k(a_0,a_1,a_{i_1},\ldots,a_{i_{k-1}}))'$$

(in terms of the leading t^m term in the formal power series)

$$\begin{aligned}
&\approx -\frac{1}{4}\alpha_1^2(t)\cdot\frac{1}{2\cdot k!}(\overrightarrow{a_0a_1}^2)'\sum_{2\le i_1<\cdots<i_{k-1}} \alpha_{i_1}(t)\cdots\alpha_{i_{k-1}}(t)\,d(a_0,a_1;a_{i_1},\ldots,a_{i_{k-1}})\\
&\approx -\frac{1}{4}\alpha_1^2(t)\cdot\frac{1}{2\cdot k!}(\overrightarrow{a_0a_1}^2)'\cdot c_{k-1}\\
&\le 0
\end{aligned}$$

The equality holds if and only if $A_0(t) = A_1(t)$, namely, $A_1(t)$, …, $A_{n+2}(t)$ stay in a common S^n (or H^n). □

Now we have the following main theorem.

Theorem 4.4 *Suppose $\mathbf{A}(t)$ moves in S^{n+1} (or H^{n+1}) and is real analytic.*

(1) If $c_{k-1} > 0$, then $G_{n,k}$ is k-unyielding in S^{n+1} (or H^{n+1}), and points $A_1(t)$, …, $A_{n+2}(t)$ will stay in a common S^n (or H^n) for small $t \ge 0$.

(2) If $c_{k-1} < 0$, then $F_{n,k}$ is k-unyielding in S^{n+1} (or H^{n+1}), and points $A_1(t)$, …, $A_{n+2}(t)$ will stay in a common S^n (or H^n) for small $t \ge 0$.

Proof. We only need to prove (1), as (2) can be proved the same way. Suppose $c_{k-1} > 0$. If points $A_1(t), \ldots, A_{n+2}(t)$ do not stay in a common S^n (or H^n), then $A_0(t) \neq A_1(t)$. By Lemma 4.4, we have

$$\sum_{1 \le i_1 < \cdots < i_{k+1}} \alpha_{i_1} \cdots \alpha_{i_{k+1}} \, |\overrightarrow{OA_{i_1}(t) \cdots A_{i_{k+1}}(t)}| \, (V_k(A_{i_1}(t), \ldots, A_{i_{k+1}}(t)))' < 0.$$

However, for $G_{n,k}$, by definition we have

$$\alpha_{i_1} \cdots \alpha_{i_{k+1}} (V_k(A_{i_1}(t), \ldots, A_{i_{k+1}}(t)))' \ge 0,$$

which is a contradiction.

As $A_1(t), \ldots, A_{n+2}(t)$ stay in a common S^n (or H^n), by applying Theorem 4.3, we then have that $G_{n,k}$ is k-unyielding in S^{n+1} (or H^{n+1}). □

Remark. We believe that Theorem 4.4 can be proved without the restriction that "$\mathbf{A}(t)$ is real analytic".

4.4 Characteristic polynomial in S^n_+ and H^n

Like we did in Euclidean space, the concept of *characteristic polynomial* can be defined on points in S^n_+ and H^n. Let $A_1, \ldots, A_m$ be m ($m \ge n+2$) points in S^n_+ or H^n, and $\alpha_1, \ldots, \alpha_m$ be a sequence of coefficients that satisfy $\sum_{i=1}^m \alpha_i A_i = 0$.

For $0 \le k \le n$, let c_k be the form in formula (4.3), we then define the characteristic polynomial of (A_i, α_i) $1 \le i \le m$ to be

$$f(x) = c_0 x^n - c_1 x^{n-1} + \cdots + (-1)^i c_i x^{n-i} + \cdots + (-1)^n c_n.$$

Conjecture 4.1 *Suppose $f(x)$ is defined in H^n.*

(1) All roots of $f(x)$ are real numbers.

(2) If $m = n+2$ and the $n+2$ points are in general positions in H^n, and there are s positive numbers in all α_i, then $f(x)$ has $s-1$ positive and $n+1-s$ negative roots.

4.5 2-tensegrity frameworks $G_{n,2}$ and $F_{n,2}$ in S^n_+ and H^n

For $G_{n,2}$ and $F_{n,2}$, the geometric meaning of $c_1 = 0$ is very simple.

Corollary 4.1 *For $G_{n,2}$ and $F_{n,2}$ in S^n_+ (or H^n), $c_1 = 0$ if and only if points $A_1, \ldots, A_{n+2}$ are in a n-dim hyperplane in R^{n+1}(or $R^{n,1}$).*

Proof. By Lemma 4.2, we have $c_1 = 2\sum_{i=1}^{n+2} \alpha_i$ for the S^n_+ case, and $c_1 = -2\sum_{i=1}^{n+2} \alpha_i$ for the H^n case. Since $\sum_{i=1}^{n+2} \alpha_i A_i = 0$, so $c_1 = 0$ if and only if points $A_1, \ldots, A_{n+2}$ are in a n-dim hyperplane. □

In Figure 3, on S^2_+, A_1 is the point that outside the dotted circle, and $c_1 > 0$. In Figure 4, on H^2, A_1 is the point that outside the dotted loop, and $c_1 > 0$. It will be interesting to know the geometric meaning of $c_k = 0$ when $k \ge 2$.

We now give a detailed discussion of $d(B, C; A)$ in H^n.



Figure 3:



Figure 4:

Proposition 4.1 *(1) For the S^n_+ case, we have*

$$d(B,C;A) = \frac{2}{1+\cos \widetilde{BC}} \overrightarrow{BA} \cdot \overrightarrow{CA}$$

(2) For the H^n case, we have

$$d(B,C;A) = \frac{2}{1+\cosh \widetilde{BC}} \overrightarrow{BA} \cdot \overrightarrow{CA}$$

Proof. Using Lemma 4.1. □

Proposition 4.1 is the analogue of Lemma 2.1, which says that $\partial_{\overrightarrow{BC}^2} \overrightarrow{ABC}^2$ is the inner product of $\overrightarrow{BA}$ and $\overrightarrow{CA}$. Indeed, we will show that in H^n, $d(B,C;A)$ does show properties that make it look like an inner product.

Given a point A in H^n, define a mapping f from H^n to the smallest linear space H^n_A associated with a bilinear product "·", such that $f(A)$ is the origin in H^n_A and

$$d(B,C;A) = f(B) \cdot f(C).$$

Theorem 4.5 *If $n = 1$, then the bilinear product "·" is positive definite in H^1_A, and H^1_A is an infinite dimensional space.*

Proof. We will prove it by showing that for any k points $P_1, \ldots, P_k$ in H^1, we have

$$\det(f(P_i) \cdot f(P_j))_{1\le i,j\le k} \ge 0.$$

Pick up a direction on H^1, we use r_i to denote $\widetilde{P_iA}$ if P_i is at the "right" side of A, and denote $-\widetilde{P_iA}$ if P_i is at the "left" side of A. Then we have

$$\det\left(f(P_i) \cdot f(P_j)\right)_{1\le i,j\le k}$$

(by Proposition 4.1)

$$\begin{aligned}
&= \det\left(\frac{2}{1+\cosh \widetilde{P_iP_j}} \overrightarrow{P_iA} \cdot \overrightarrow{P_jA}\right) \\
&= \det\left(\frac{2}{1+\cosh(r_i - r_j)} (\cosh r_i + \cosh r_j - \cosh(r_i - r_j) - 1)\right) \\
&= \det\left(\frac{4}{\cosh\frac{r_i - r_j}{2}} \sinh\frac{r_i}{2} \sinh\frac{r_j}{2}\right) \\
&= 2^k \left(\prod_i (e^{r_i} - 1)\right)^2 \det\left(\frac{1}{e^{r_i} + e^{r_j}}\right)_{1\le i,j\le k} \\
&\ge 0.
\end{aligned}$$

The last step uses Cauchy's determinant identity [3], which says that

$$\det\left(\frac{1}{x_i + y_j}\right)_{1\le i,j\le n} = \frac{\prod_{i<j}(x_j - x_i)(y_j - y_i)}{\prod_{i,j}(x_i + y_j)}.$$

The equality of the last step holds if and only if $P_i = A$ for some i or $P_i = P_j$ for some i and j, which implies that H^1_A is an infinite dimensional inner product space. □

Conjecture 4.2 *For any $n > 1$, the bilinear product "$\cdot$" is positive definite in H^n_A.*

4.6 A special version of k-tensegrity Frameworks $G_{n,k}$ and $F_{n,k}$ in S^n

In S^n, we now briefly discuss a special case that when the points $A_1, \ldots, A_{n+2}$ are not in a hemisphere. Suppose $A_1, \ldots, A_{n+2}$ are in general position in S^n, and the origin O is inside the *Euclidean* $(n+1)$-simplex $A_1 \cdots A_{n+2}$. We let $G_{n,k}$ be a k-tensegrity framework whose k-faces are all k-struts; and $F_{n,k}$ be the one whose k-faces are all k-cables.

Theorem 4.6 *(1) If $\mathbf{A}(t)$ is real analytic, then $G_{n,k}$ is k-unyielding in S^{n+1}.*

(2) If $\mathbf{A}(t)$ is C^1, then $F_{n,k}$ is k-unyielding in S^n.

Proof. Can be proved by slightly modifying Theorem 4.4 and Theorem 4.3 respectively. □

E-mail address: lizhaozhang@alum.mit.edu

References

[1] K. Bezdek and R. Connelly, *Two-distance preserving functions from Euclidean space*, Discrete geometry and rigidity(Budapest, 1999), Period. Math. Hungar., **39** (1999), 185-200.

[2] A.L. Cauchy, *Sur les polygones et les polyèdres*, J. École Polytechnique, XVIe Cahier, Tome IX, 87-98.

[3] A.L. Cauchy, *Mémoire sur les fonctions alternées et sur les sommes alternées*, Exercices Anal. et Phys. Math. **2** (1841), 151-159.

[4] R. Connelly, *Rigidity*, in: Handbook of convex geometry, Vol. A, Amesterdam; New York: North-Holland, 1993, pp. 223–271.

[5] R. Connelly, *A counterexample to the rigidity conjecture for polyhedra*, Inst. Hautes Études Sci. Publ. Math., **47**, (1978), 333-338.

[6] R. Connelly, I. Sabitov and A. Walz, *The bellows conjecture*, Contributions to Algebra and Geometry, **38** (1997), no.1, 1-10.

[7] W. Fenchel, Elementary Geometry in Hyperbolic Space, Berlin; New York: W. de Gruyter, 1989.

[8] J. Milnor, *The Schläfli differential equality*, Collected papers, Vol. 1, Publish or Perish, 1994.

[9] I. Sabitov, *On the problem of invariance of the volume of a flexible polyhedron*, Russian Mathematical Surveys, **50** (1995), no.2, 451-452.

[10] T.-S. Tay, N. White and W. Whiteley, *Skeletal rigidity of simplicial complexes, I*, European J. Combinatorics, **16** (1995), 381-403.

[11] T.-S. Tay, N. White and W. Whiteley, *Skeletal rigidity of simplicial complexes, II*, European J. Combinatorics, **16** (1995), 503-523.

[12] L. Zhang, Rigidity and Invariance Properties of Certain Geometric Frameworks, PhD Thesis, MIT (2002) [available from: http://dspace.mit.edu/handle/1721.1/8337]